

# Abstract

Prostaglandins Leukot Essent Fatty Acids. 2010 Sep;83(3):137-41.

## Bioavailability of marine n-3 fatty acid formulations.

Dyerberg J, Madsen P, Møller JM, Aardestrup I, Schmidt EB.

Department of Human Nutrition, Faculty of Life Sciences, University of Copenhagen, Copenhagen, Denmark.

**BACKGROUND:** The use of marine n-3 polyunsaturated fatty acids (n-3 PUFA) as supplements has prompted the development of concentrated formulations to overcome compliance problems.

**OBJECTIVE:** The present study compares three concentrated preparations - ethyl esters, free fatty acids and re-esterified triglycerides - with placebo oil in a double-blinded design, and with fish body oil and cod liver oil in single-blinded arms.

**METHODS:** Seventy-two volunteers were given approximately 3.3g of eicosapentaenoic acid (EPA) plus docosahexaenoic acid (DHA) daily for 2 weeks. Increases in absolute amounts of EPA and DHA in fasting serum triglycerides, cholesterol esters and phospholipids were examined.

**RESULTS:** Bioavailability of EPA+DHA from re-esterified triglycerides was superior (124%) compared with natural fish oil, whereas the bioavailability from ethyl esters was inferior (73%). Free fatty acid bioavailability (91%) did not differ significantly from natural triglycerides.

**CONCLUSIONS:** The stereochemistry of fatty acid in acylglycerols did not influence the bioavailability of EPA and DHA.

PMID: 20638827